

Holographic Limitations of the Effective Field Theory of Inflation

Andreas Albrecht, Nemanja Kaloper and Yong-Seon Song
 Department of Physics, University of California, Davis, CA 95616
 (February 1, 2008)

Holographic considerations may provide a glimpse of quantum gravity beyond what is currently accessible by other means. Here we apply holography to inflationary cosmology. We argue that the appropriate holographic bound on the total entropy in the inflationary perturbations is given by the *difference* of the area of the apparent horizon in the slow-roll approximation and its exact value near the end of inflation. This implies that the effective field theory of the inflaton weakly coupled to gravity requires modifications below the Planck scale. For a conventional model of inflation this scale is of order of the GUT scale, $\Lambda_{UV} \lesssim 10^{16} \text{GeV}$, but could be considerably lower. Signatures of such new physics could show up in the CMB.

We live in a big, old, but nearly wrinkle-free universe. Inflation is currently the best paradigm which could explain how it got its face-lift [1]. However, a truly fundamental understanding of inflation is yet to emerge from yet to be discovered complete theory of quantum gravity. We are therefore far from building cosmology directly from a complete quantum gravity theory (see, e.g. [2]-[8]). Nevertheless, there have been remarkable developments in black hole physics which led to the discovery of general principles, originating from black hole thermodynamics, that appear to be universal and transcend any given specific approach to quantum gravity. 't Hooft [9] and Susskind [10] have named this set of rules the holographic principle, showing that for black holes, it is *de facto* the solution of the black hole information problem. It states that a black hole has far fewer degrees of freedom than expected, with its total entropy given by the quarter of the horizon area in Planck units, as per Bekenstein-Hawking formula [11,12]. It has since been shown that generic field theories coupled to gravity overcount the degrees of freedom, by deducing “holographic area bounds” which limit the number of physical degrees of freedom. In this Letter we explore the implications of these ideas for inflation.

Fischler and Susskind have made the first step towards incorporating the holographic principle in cosmology [13], suggesting that the entropy which crosses the lightlike boundary of an observable region of the universe, the particle horizon, should not exceed the horizon area in Planck units. Subsequent to [13,14], Bousso has offered a covariant version of the holographic bound [15], outlining a prescription for choosing the holographic “screens”. The idea is to take any lightlike surface ending on either a caustic or the apparent horizon, and limit the entropy which crosses it by a 1/4 of the maximal area along the light sheet in Planck units. This seems to produce desirable results for matter with generic properties [16]. The implications of holography for cosmology have been explored in many works (e.g. [6]-[8], [3], [17], [18]).

Here we argue that the holographic principle imposes a limit on the validity of the description of inflation, and specifically inflationary fluctuations, via an effective

field theory weakly coupled to gravity in the causal patch (EFTwg). The crux of our argument is that the entropy deposited into the metric fluctuations generated by the inflaton could easily exceed the holographic bounds, if the EFTwg is valid to arbitrarily short distances. This cannot happen in a truly holographic theory. An example is a field theory in a black hole background, where the entropy, defined as the logarithm of the partition function, in a causal region of a black hole spacetime, is badly divergent [19]. Only after renormalizing it does one recover the Bekenstein-Hawking area law [19]. This corresponds to cutting off the modes in the UV, residing close to the horizon. We argue that in inflation this new physics encoded in the UV cutoff affects the inflationary fluctuations, which are classical scalar modes with their wavelengths greater than the instantaneous Hubble length. These modes start out as short distance excitations inside the Hubble horizon before they are expelled out by the superluminal expansion, and hence are sensitive to the physics represented by the UV cutoff.

In the absence of a precise formulation we can crudely approximate the holographic redundancies by an energy cutoff Λ_{UV} on EFTwg, which remains sensitive to the dynamics. This cutoff is a reflection of the expected strong gravitational interactions close to the horizon. We find that it is bounded from above, and the bound is inversely proportional to the third root of the number of e-folds. This implies that the number of e-folds which can be described by EFTwg in the causal patch is limited. In the commonly quoted example of inflation occurring when $H \lesssim 10^{14} \text{GeV}$, and lasting about ~ 100 e-folds, the cutoff is $\Lambda_{UV} \lesssim 10^{16} \text{GeV} \sim M_{GUT}$, but it could be lower depending on the model of inflation. A similar scale also appears as the upper bound on the scale of new physics which might imprint an observable signature in the CMB as the $\mathcal{O}(H^2/\Lambda_{UV}^2)$ corrections on top of the leading order fluctuations [20]. We submit that these signatures could be a reflection of holography, although a deeper knowledge of the underlying physics is required to determine their detailed form.

We start with the picture of pure de Sitter space as a “hot tin can” [3], with an observer in the causal patch

(the interior of the “can”) surrounded by a hot horizon (the “walls”). The physical degrees of freedom, accessible to the observer, are in thermal equilibrium with the walls, at the Gibbons-Hawking temperature. In the case of nonzero, constant slow roll parameters, such as in a quintessence-dominated universe, the picture of [7] suggested that the ensuing causal region can be viewed as a hot tin can with moving walls, so that the temperature of the gas in equilibrium inside the cavity decreases due to the expansion of the can. An inflating region in the slow roll regime closely resembles de Sitter space, with a key difference: unlike the perfect de Sitter space, where the apparent horizon and the event horizon coincide and so there is no “exterior” of the can, during inflation modes in the IR are expelled from the apparent horizon. These modes are real physical degrees of freedom, and may or may not ever thermalize depending on the future evolution of the universe. We will divide the total instantaneous entropy in the inflating patch into the dominant equilibrium contribution, approximated by the horizon area of the would-be pure de Sitter space, and a small non-equilibrium contribution, where the inflaton fluctuations count in. We will argue that this is analogous with the familiar case of a black hole in de Sitter space. In our picture inflation is a hot, expanding, but porous tin can, slowly leaking entropy into a cold reservoir via the expelled modes.

Assuming the standard theory of inflationary fluctuations below the cutoff Λ_{UV} , we estimate the entropy-per-mode by counting the mode’s quantum excitation level [21,22]. We argue that this entropy is bounded by the *difference* of the areas of true apparent horizon during inflation, and its slow-roll value. This goes beyond the covariant holographic bounds [15] in a crucial way, because it allows us to discriminate between the entropy of modes in equilibrium and the entropy in density fluctuations which froze out. Our bounds constrain Λ_{UV} to physically interesting values, which depend on the inflation model, but in general are significantly below M_p .

We will work with the flat 4D FRW metric and ignore any non-inflaton matter sources (both good approximations during inflation). The background obeys

$$3H^2 = \frac{\dot{\phi}^2}{2M_p^2} + \frac{V(\phi)}{M_p^2}, \quad \ddot{\phi} + 3H\dot{\phi} + \frac{\partial V(\phi)}{\partial \phi} = 0, \quad (1)$$

where the overdot is the time derivative, $H = \dot{a}/a$ is the Hubble parameter, a is the FRW scale factor, ϕ is the inflaton, and $V(\phi)$ is the inflaton potential. The Planck mass is defined by $M_p = 1/\sqrt{8\pi G_N}$ since $c = \hbar = 1$.

Inflation occurs when the potential energy of the scalar dominates in (1). The dynamics is well approximated by the slow-roll regime, formally defined by the standard conditions $|\epsilon|, |\delta| \ll 1$ where the slow-roll parameters are $\epsilon = \frac{3\dot{\phi}^2}{2V}$, $\delta = \frac{\ddot{\phi}}{H\dot{\phi}}$. The slow-roll approximation then consists of dropping the terms of order $\mathcal{O}(\epsilon, \delta)$. Then (1) reduce to

$$3H^2 = \frac{V}{M_p^2}, \quad 3H\dot{\phi} + \frac{\partial V}{\partial \phi} = 0. \quad (2)$$

We stress here that in the slow-roll approximation the potential, and thus the radius of the apparent horizon (i.e. the Hubble horizon), defined as the hypersurface at $l_{AH} = H^{-1} = M_p/\sqrt{3V}$ from the origin, is essentially constant during each Hubble time. Although V is implicitly time-dependent, as seen from Eq. (2), it changes during a single efold by at most $\Delta V \lesssim |\frac{V'\Delta\phi}{V}| \simeq \frac{\epsilon}{2}V$. Because one *defines* the slow-roll by dropping $\mathcal{O}(\epsilon, \delta)$ terms, the solutions are only reliable to this order, and so the radius of the apparent horizon is constant with the error of only $\mathcal{O}(\epsilon)$. This is crucial for our arguments to follow.




FIG. 1. Penrose diagram of an inflationary spacetime.

The causal structure of the inflating region is given in Fig. 1 [20]. The future null line is the future null causal boundary (FNCB), which could be either a null infinity, or an event horizon if the universe ends up inflating forever. The past null line is the particle horizon (PH), measuring how far the light travelled since the birth of the universe (taken as $t = 0$), given by $L_H(t) = a(t) \int_0^t \frac{dt'}{a(t')}$. The shaded area (R) represents reheating, with inflation occurring before it, and the radiation/matter era after it. The apparent, i.e. Hubble, horizon (AH) is the bold solid curve, during inflation almost null and directed first outward, turning inward later. This is because $H \approx \text{const.}$ during inflation, and reflects the thermal nature of the quantum state during inflation [20], an approximation valid over the time scales of order the Hubble time as argued above. According to the complementarity principle, all active experiences of an observer can be fully described by the physics in the causal patch, or the tin can [6,3,4,7]. As we are only interested in the inflationary fluctuations pushed in between AH and FNCB, we will ignore the exterior of the causal patch as immaterial. LS is a light-sheet defining a holographic screen.

The out-of-equilibrium modes originate from within the apparent horizon, and are expelled out of it by inflationary expansion. They remain frozen until they cross back into the apparent horizon after inflation, where they grow due to the gravitational instability. The entropy per mode is found by considering the fluctuations as squeezed states of the inflaton [21,22]. One should sum

up the entropy in all modes which were expelled out of the apparent horizon during inflation, with momenta determined by the horizon crossing matching condition and the full range of the Hubble scale during inflation where EFTwg is valid, $H(t) \in [H, \Lambda_{UV}]$. The key point is that the result is UV-dominated, since the inflationary fluctuations are nearly scale-invariant, with $P_\Phi(p) \sim k^{n-1}$ where $n = 1 + \mathcal{O}(\epsilon, \delta)$ is close to unity as $|\epsilon|, |\delta| \ll 1$, and so we can *i)* approximate the entropy per mode by an (almost) constant value, of order unity and *ii)* assume that the modes are pushed out of the horizon at a rate independent of their wavelength. The entropy deposited in them will only depend on the number of modes pushed out of the horizon. The modes which left the horizon at the beginning will be the most numerous, since at that time the Hubble length was the shortest possible, and they will dominate the entropy. Because their wavelength was of order of the Hubble length in the beginning of the EFTwg description of inflation, $\lambda_{UV} \simeq \Lambda_{UV}^{-1}$, their number per horizon volume at the end of inflation is of order of the number of the initial Hubble domains inside the thermal patch at the end of inflation, $n \simeq 4\pi(\lambda_{thermal}/\lambda_{UV})^3 \simeq 4\pi(\Lambda_{UV}/H)^3$. Thus, the entropy in the expelled modes is

$$S \simeq 4\pi \frac{\Lambda_{UV}^3}{H^3}. \quad (3)$$

The calculations of [21,22] bear out this estimate. Eq. (3) presumes that fluctuations leave the apparent horizon, which is only possible if it is timelike, so that the slow roll parameters are small but not exactly zero. To take the limit $\epsilon, \delta \rightarrow 0$ one must take $\Lambda_{UV} \rightarrow H$ and keep the subleading corrections to (3), where we expect $S \rightarrow 0$.

The holographic principle gives us another way of estimating the entropy in the inflating patch. A rule for bounding the entropy is to pick a light-sheet crossed by the entropy, and limit the entropy by the maximal area along the light-sheet [13,15]. A suitable light-sheet is given by the faint solid null line (LS) in Fig. 1, which is the backward light cone of an observer near the end of inflation, with maximal area $\mathcal{A} \simeq M_p^2/H^2$. We chose it so that it maximizes the bounds, as it intersects the apparent horizon as close as possible to the end of inflation, where the slow-roll conditions still hold, and so it is crossed by (almost) all of the modes expelled out of the apparent horizon. However, the ensuing bounds still are not very strong [13], because they do not discriminate between the modes in equilibrium and those out of it.

We propose here that the bounds could be strengthened by separating these modes from each other during inflation. To motivate the procedure we first consider black holes in de Sitter space [23]. They are the most stable objects in a background de Sitter space: they can be viewed as small, hot, balls of gas in a cold cavity, which are very slowly melting away by emitting Hawking radiation. Because this process is very slow, given by the time $\tau_{BH} \simeq M_{BH}^3/M_p^4$, most of the energy which slipped

into a black hole will not thermalize with de Sitter background for a very long time. Their entropy is given by their horizon area, which for black holes smaller than the de Sitter radius is as in the flat space, $S_{BH} \sim M_p^2 \mathcal{A}_{BH} \sim 4\pi M_p^2 r_{BH}^2 \sim (M_{BH}/M_p)^2$. On the other hand, the equilibrium entropy in the asymptotically de Sitter geometry is approximately $S_{eq} \sim 4\pi M_p^4/\lambda - \mathcal{O}(1)M_p^2 \sqrt{\mathcal{A}_{dS} \mathcal{A}_{BH}}$. Thus the entropy in the black hole, in modes out of equilibrium with de Sitter cavity, is smaller than the *difference* between the entropy of the modes in equilibrium and the total entropy in pure de Sitter space with the same value of the cosmological constant:

$$S_{BH} \lesssim \sqrt{S_{BH} S_{eq}} \lesssim S_{dS} - S_{eq}. \quad (4)$$

We will now adopt a similar formula for the case of inflation. If the slow-roll parameters were exactly zero, the apparent horizon would have coincided with the event horizon, and nothing would ever leave it (or freeze out) in a finite observer's time. Thus in this limit the entropy in the metric fluctuations would vanish too. Slow time-dependence of H is thus essential for producing the fluctuations. However, in the slow-roll approximation during the times $\lesssim H^{-1}$ we have seen above that the Hubble horizon is practically indistinguishable from a *true* event horizon, since $\Delta V = 0$ to leading order in ϵ . Hence the slow-roll Hubble horizon has no ability to distinguish the modes in and out of equilibrium. Its area in the slow-roll regime encodes the contribution from all the field theory modes generated in the Hubble patch. In contrast, the true apparent horizon discriminates between the modes in equilibrium, with wavelengths shorter than the Hubble length, and the modes pushed out of equilibrium, which grow greater than the Hubble patch. In fact, the apparent horizon moves inward toward the center of the inflating region relative to the slow roll Hubble horizon, as does the cosmological horizon in Schwarzschild-de Sitter space-time, to record that there is fewer modes in equilibrium. In a manner of speaking, if exact de Sitter space is a cavity with hot static walls, the inflating patch is a cavity with hot, moving and porous walls, which lets out some of the IR degrees of freedom. To extract their entropy, we take the difference of the slow-roll and exact areas of the apparent horizon,

$$\Delta \mathcal{A} = \frac{\epsilon}{3} \mathcal{A}. \quad (5)$$

To leading order in the slow-roll parameters we can take $\mathcal{A}$ on the RHS to be the slow roll horizon area, or equivalently the horizon area of the would-be de Sitter space found in the limit $\epsilon, \delta \rightarrow 0$, since the errors are of order $\mathcal{O}(\epsilon^2) \ll 1$. Then the field theory result for the entropy cannot exceed this area in Planck units,

$$S \lesssim \frac{\epsilon}{3} M_p^2 \mathcal{A}. \quad (6)$$

This bound on the entropy in inflationary perturbations is stronger than the covariant, all-inclusive, bound of [15]

by the power of $\epsilon \ll 1$. Using (3), we rewrite this as the bound on the UV cutoff Λ_{UV} :

$$\Lambda_{UV} \lesssim \left(\frac{\epsilon}{3}\right)^{1/3} (HM_p^2)^{1/3}. \quad (7)$$

This is our main result. It tells us that the causal patch description of inflation by the effective field theory weakly coupled to gravity cannot be valid down to arbitrarily short distances. Instead, the holographic redundancies must cut off the entropy that can fit in the apparent horizon, including the portion in the IR which can get pushed out of equilibrium. Let us consider a simple explicit example applying (7) to chaotic inflation [24]. A typical chaotic inflation potential is of the form $\lambda\phi^n/n$ for an arbitrary (integer) power n . The slow roll parameter ϵ is related to the number of efolds before the end of inflation $\mathcal{N}$ by $\epsilon = \frac{n}{4\mathcal{N}}$, so we can rewrite (7) as

$$\Lambda_{UV} \lesssim \left(\frac{n}{12\mathcal{N}}\right)^{1/3} (HM_p^2)^{1/3}. \quad (8)$$

Thus, with $H \lesssim 10^{14} \text{ GeV}$ and $\mathcal{N} \sim 100$, we get

$$\Lambda_{UV} \lesssim 10^{16} \text{ GeV}, \quad (9)$$

so EFTwg receives modifications below the GUT scale. A longer inflation requires a lower cutoff. Note that (7)-(9) are approximations, valid as long as $\Lambda_{UV} \gg H$ and $1 \gg \epsilon > 0$, and that they are a bound on the semiclassical regime of inflation. It would be interesting to see what is the relation between eternal inflation and the causal patch description [25].

Our conclusion is that the holographic principle limits the validity of the causal patch EFTwg description of the inflaton, implying modifications at a scale set by (7), below M_p . We do not necessarily view this as a *conceptual* obstruction to inflation. We see our result as a hint that the inflaton need not be an elementary field, but emerges from the fundamental theory merely as an order parameter. At energy scales above the cutoff it would contain too much entropy, and so its EFTwg description cannot remain valid to such high energies. Alternatively, it cannot model an arbitrarily long period of inflation. An interesting consequence of our arguments is related to the recent field theory analyses of the signatures of high energy physics in the CMB. There scales similar to our cutoff arose as the upper bounds on scales of the new physics which might imprint subleading signatures on the CMB [20]. Because we expect the usual effective field theory to hold at scales $H \lesssim 10^{14} \text{ GeV} < \Lambda_{UV}$ (so that we have an effective field theory model of inflation in the first place), the effects of the holographic redundancies could be similar to the signatures of new physics near the cutoff Λ_{UV} . In the EFTwg regime the corrections to the leading order results appear as $\mathcal{O}(H^2/\Lambda_{UV}^2)$ effects [20]. If the new physics is below the GUT scale the corrections could beat the cosmic variance, and may be an exciting prospect for the observational signatures of holography in the sky.

We thank T. Banks, W. Fischler, M. Kaplinghat, M. Kleban, L. Kofman, A. Lawrence, A. Linde, S. Shenker and L. Susskind for valuable discussions. A.A. and Y-S.S. are supported by DOE grant DE-FG03-91ER40674. N.K. was supported by the UCD, by NSF Grant PHY-9870115 to the Stanford ITP, and by NSF Grant PHY-99-07949 to the UCSB KITP.

-
- [1] A. Guth, Phys. Rev. D **23**, 347 (1981); A. Linde, Phys. Lett. B **108**, 389 (1982); A. Albrecht and P. Steinhardt, Phys. Rev. Lett. **48**, 1220 (1982).
 - [2] A. Linde, Phys. Lett. B **162** (1985) 281; A. Albrecht et al, Phys. Rev. D **35**, 429 (1987); E. Farhi et al, Phys. Lett. B **183**, 149 (1987); D. Goldwirth et al, Phys. Rept. **214**, 223 (1992); A. Borde et al, Phys. Rev. Lett. **72**, 3305 (1994); A. Borde et al, gr-qc/0110012; S. Hollands et al, gr-qc/0205058; A. Albrecht, astro-ph/0210527.
 - [3] L. Dyson, M. Kleban and L. Susskind, hep-th/0208013.
 - [4] L. Dyson, J. Lindesay and L. Susskind, hep-th/0202163.
 - [5] T. Banks et al, JHEP **9901**, 019 (1999).
 - [6] T. Banks, hep-th/0007146.
 - [7] S. Hellerman et al, JHEP **0106**, 003 (2001); W. Fischler et al, JHEP **0107**, 003 (2001).
 - [8] E. Witten, hep-th/0106109.
 - [9] G. 't Hooft, gr-qc/9310026.
 - [10] L. Susskind, J. Math. Phys. **36**, 6377 (1995).
 - [11] J. Bekenstein, Phys. Rev. D **7** (1973) 2333; Phys. Rev. D **9**, 3292 (1974).
 - [12] S. Hawking, Nature **248** (1974) 30; Commun. Math. Phys. **43** (1975) 199.
 - [13] W. Fischler and L. Susskind, hep-th/9806039.
 - [14] R. Easther et al, Phys. Rev. Lett. **82**, 4967 (1999); D. Bak et al, Class. Quant. Grav. **17**, L83 (2000); G. Veneziano, Phys. Lett. B **454**, 22 (1999); N. Kaloper et al, Phys. Rev. D **60**, 103509 (1999).
 - [15] R. Bousso, JHEP **9907**, 004 (1999); JHEP **9906**, 028 (1999).
 - [16] E. Flanagan et al, Phys. Rev. D **62**, 084035 (2000).
 - [17] T. Banks et al, hep-th/0102077; hep-th/0111142.
 - [18] C. Hogan, Phys. Rev. D **66**, 023521 (2002).
 - [19] G. 't Hooft, Nucl. Phys. B **256**, 727 (1985).
 - [20] N. Kaloper et al, hep-th/0201158; N. Kaloper et al, hep-th/0209231.
 - [21] L. Grishchuk et al, Phys. Rev. D **42**, 3413 (1990).
 - [22] R. Brandenberger et al, Phys. Rev. Lett. **69**, 3606 (1992); Phys. Rev. D **48**, 2443 (1993).
 - [23] G. Gibbons et al, Phys. Rev. D **15** (1977) 2738.
 - [24] A. Linde, Phys. Lett. B **129**, 177 (1983).
 - [25] In fact, the bound (8) suggests that in the causal patch EFTwg the vev of the inflaton must be below the self-reproduction value. It will be interesting to further explore the implications of this.